Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 7 November 2011 (MN LaTeX style file v2.2)

arXiv:1111.0978v1 [astro-ph.SR] 3 Nov 2011

Abundance analysis of Am binaries and search for tidally driven abundance anomalies - III. HD 116657, HD 138213, HD 155375, HD 159560, HD 196544 and HD 204188

I. Stateva[1]*, I.Kh. Iliev[1], J. Budaj[2]
[1] *Institute of Astronomy with NAO, Bulgarian Academy of Sciences, Sofia 1784, Bulgaria*
[2] *Astronomical Institute, Slovak Academy of Sciences, 059 60 Tatranská Lomnica, The Slovak Republic*

Accepted ; Received

ABSTRACT

We continue here the systematic abundance analysis of a sample of Am binaries in order to search for possible abundance anomalies driven by tidal interaction in these binary systems.

New CCD observations in two spectral regions (6400–6500, 6660–6760 ÅÅ) of HD 116657, HD 138213, HD 155375, HD 159560, HD 196544 and HD 204188 were obtained. Synthetic spectrum analysis was carried out and basic stellar properties, effective temperatures, gravities, projected rotational velocities, masses, ages and abundances of several elements were determined. We conclude that all six stars are Am stars.

These stars were put into the context of other Am binaries with $10 < P_{\rm orb} < 200$ days and their abundance anomalies discussed in the context of possible tidal effects. There is clear anti-correlation of the Am peculiarities with $v\sin i$. However, there seems to be also a correlation with the eccentricity and may be with the orbital period. The dependence on the temperature, age, mass, and microturbulence was studied as well. The projected rotational velocities obtained by us were compared to those of Royer et al. (2002) and Abt & Morrell (1995).

Key words: Stars: chemically peculiar – Stars: abundances – Stars: individual: HD 116657, HD 138213, HD 155375, HD 159560, HD 196544, HD 204188 – binaries: close – diffusion – hydrodynamics

1 INTRODUCTION

The Am stars is a well known subgroup of chemically peculiar (CP) stars on the upper Main Sequence (MS). They exhibit abnormally strong metallic and unusually weak Ca and Sc lines and, as a consequence, the spectral types inferred from calcium lines are usually earlier than those from hydrogen lines, and the latter are earlier than the spectral types from the metallic lines. The anomalous intensity of most of these absorption lines is due to the abnormal chemical composition of superficial layers. The typical abundance pattern of Am stars is that they exhibit a deficit of light elements like C, Mg, Ca, Sc and progressively increasing overabundances of iron group and heavier elements. This abundance pattern is often referred to as the Am phenomenon. Rotation was found to play a key role in these stars and there is a growing amount of recent observational evidence that Am peculiarity is either a smooth or a step function of rotation (Iliev & Budaj 2008 and the references herein; Abt 2000; Burkhart & Coupry 2000). Nevertheless, Am peculiarity does seem to depend on evolutionary status or age as well. It may (1) develop very quickly soon after the star arrives on the MS, or even before that (Burkhart & Coupry 2000), and does not undergo considerable changes during the MS phase, or (2) observable abundances of some elements may vary with age and this can be used to constrain the evolutionary models (Monier & Richard 2004; Monier 2005). At the same time, no significant correlation of the abundance anomalies with $v\sin i$ were found by Monier & Richard (2004), and Monier (2005). Apart from that the Am phenomenon is apparently restricted to a well-defined region of the MS in the HR diagram which implies its dependence on atmospheric parameters such as effective temperature and gravity (Künzli & North 1998; Hui-Bon-Hoa 2000).

The Am peculiarity seems to depend on the orbital elements in a binary system as well. Budaj (1996), Budaj (1997), Iliev et al. (1998) studied $v\sin i$ vs $P_{\rm orb}$, e vs $P_{\rm orb}$,

---

* E-mail: stateva@astro.bas.bg

δm_1 vs $P_{\rm orb}$, $f(m)$ vs $P_{\rm orb}$, $v\,\sin i$ vs $P_{\rm p}$, δm_1 vs $P_{\rm p}$, where $P_{\rm orb}$ is the orbital period, δm_1 is a metallicity parameter which shows the difference in the dereddened m_1 index of $uvby\beta$ photometry between an Am star and a normal star of the same index, $f(m)$ is a mass function, and P_p is the "instantaneous" orbital period at periastron. The authors concluded that there are a number of subtle effects which are difficult to understand within the current framework of the rotation and atmospheric parameters as the only agents determining the Am star peculiarity.

Since Am stars are often found in binaries (North et al. 1998; Debernardi et al. 2000) they provide a unique opportunity to study the influence of a companion on the stellar hydrodynamics.

This forced us to study the Am peculiarity and the orbital elements and $v\,\sin i$, mass, age as well. In order to explore the possible dependence of Am peculiarity on the orbital elements of the binary system we started a systematic spectroscopic investigation of Am stars. A few tens of Am binaries from Budaj (1996) were chosen to be studied in details. The following criteria were applied in order to compile the star's list: targets with the declination $\delta > -10°$ and brighter than the 7-th magnitude in V-filter. In order to cover a full range of eccentricities and avoid strong synchronization effects we have chosen only stars with orbital periods $10^d < P_{\rm orb} < 200^d$. No constraints were put on the rotational velocity.

In this connection the studies of individual Am binaries are very important (e.g., Fossati et al. 2007; Zverko et al. 2008; Zverko et al. 2009; Boffin 2010; Hubrig, González & Schöller 2010; Mikulashek et al. 2010; Quiroga, Torres & Cidale 2010). Obtaining new orbital elements of many Am binaries are also needed to prove the dependence between orbital elements and the Am peculiarity (Debernardi 2002; Carquillat & Prieur 2007; Zhao et al. 2007; Fekel & Williamson 2010).

This is the last in a series of papers (Budaj & Iliev (2003), hereafter Paper I and Iliev et al. (2006), hereafter Paper II) aiming at studying together the Am peculiarity in multi-dimension parameter space involving the orbital elements, $v\,\sin i$, mass, and age.

2 OBSERVATIONS AND SAMPLE STARS

Our spectroscopic observations were carried out with the 2-m RCC telescope of the Bulgarian National Astronomical Observatory in the frame of our scientific project on Am stars in binary systems. As in the previos papers of the series we observed each star in two spectral regions: 6400–6500 ÅÅ (Ca) and 6660–6760 ÅÅ (Li). For more details about the processing procedures see these papers.

The focus of this paper is to continue the analysis of another six stars from the sample, namely HD 116657, HD 138213, HD 155375, HD 159560, HD 196544 and HD 204188.

Table 1. Log of observations: Spectrum number, date [dd.mm.yyyy], HJD (2450000+) of the beginning of the exposure, effective exposure time [min], spectral region, heliocentric radial velocity of the primary and its error [km s^{-1}].

N	Date	HJD	Exp.	Reg.	RV1	ΔRV1
			HD 116657			
1	04.01.2001	1913.554	40	Ca	-11.1	2.6
2	04.01.2001	1913.594	40	Li	-11.1	2.8
			HD 138213			
1	10.06.2001	2071.325	45	Ca	-12.4	1.7
2	09.06.2001	2070.357	45	Li	-11.2	3.1
			HD 155375			
1	10.06.2001	2071.392	90	Ca	24.7	2.8
2	21.08.2001	2508.300	115	Ca	19.1	1.9
3	26.08.2002	2513.251	90	Ca	25.6	1.8
4	28.08.2002	2515.292	40	Ca	24.7	1.5
5	22.09.2007	4366.270	60	Ca	8.4	2.8
6	23.09.2007	4367.245	90	Ca	10.7	1.9
7	24.09.2007	4368.229	60	Ca	14.9	2.0
8	09.06.2001	2070.430	75	Li	24.6	5.3
			HD 159560			
1	10.06.2001	2071.453	40	Ca	-23.7	2.5
2	23.08.2007	4336.395	25	Ca	-13.8	2.2
3	25.08.2007	4338.442	20	Ca	-12.7	2.9
4	11.06.2001	2072.384	45	Li	-24.6	2.7
			HD 196544			
1	10.06.2001	2069.546	60	Ca	-29.9	2.5
2	28.08.2001	2150.278	100	Ca	-7.2	3.0
3	02.09.2001	2155.332	30	Ca	12.6	4.7
4	18.05.2002	2412.560	40	Ca	-19.9	2.4
5	10.08.2000	1767.324	115	Li	21.1	4.1
			HD 204188			
1	23.07.2000	1749.437	180	Ca	-18.5	2.5
2	22.07.2000	1748.454	260	Li	-8.0	2.9



Figure 1. The location of our six stars in the HR diagram. Evolutionary tracks for $M = 3.0, 2.5, 2.0, 1.7, 1.5,$ and $1.25\ M_\odot$ are shown by solid lines, an isochrone for $\log T = 3.0$ in [yr] (ZAMS) is shown by the dotted line (Lejeune & Schaerer 2001).

Table 2. Photometry, atmospheric parameters and other relevant information about the observed stars

Star	HD 116657	HD 138213	HD 155375	HD 159560	HD 196544	HD 204188
UBV photometry						
V	2.227^T	6.146	6.586	4.865	5.433	6.078
B-V	0.057^T	0.10	0.085	0.279	0.052	0.22
U-B	–	0.12	0.086	0.068	0.039	0.06
$uvby\beta$ photometry						
E(b-y)	-0.010	-0.014	-0.005	0.001	-0.010	0.000
$(b-y)_0$	0.063	0.046	0.047	0.175	0.022	0.142
m_0	0.239	0.191	0.197	0.208	0.186	0.199
c_0	0.911	1.141	1.025	0.747	1.014	0.777
β	2.886	2.860	2.885	2.772	2.911	2.806
$T_{\rm eff}$	8470	8430	8560	7460	9090	7780
$\log g$	4.32	3.61	4.09	4.14	4.33	4.30
Geneva photometry						
U	–	1.648	1.561	1.478	1.501	1.434
V	–	0.825	0.842	0.609	0.905	0.677
B1	–	0.909	0.910	0.971	0.902	0.953
B2	–	1.445	1.449	1.404	1.469	1.422
V1	–	1.524	1.539	1.334	1.601	1.390
G	–	1.989	2.000	1.732	2.084	1.807
$T_{\rm eff}$	–	8403	8657	7321	9301	-
$\log g$	–	3.65	4.02	4.26	4.29	-
$P_{\rm orb}$	175.6[1]	105.95[2]	23.2[3]	38.0[4]	11.0[5]	21.72[5]
e	0.46[1]	0.0[2]	0.42[3]	0.03[4]	0.23[5]	0.0[6]
π	41.73 ± 0.61	6.37 ± 0.29	10.13 ± 0.45	32.80 ± 0.18	17.26 ± 0.33	21.57 ± 0.56
M_V	0.33	0.17	1.61	2.44	1.62	2.75
adopted atmospheric parameters						
$T_{\rm eff}$	8470	8500	8610	7390	9200	7780
$\log g$	4.32	3.50	4.06	4.20	4.31	4.30

Note: [1] – Gutmann (1965); [2] – Lucy & Sweeney (1971); [3] – Debernardi (2002); [4] – Margoni et al. (1992); [5] – Harper (1935); [6] – Batten, Fletcher & Mann (1978); [T] – Hipparcos and Tycho catalogue (ESA 1997); $T_{\rm eff}$ is in [K], $\log g$ in CGS units, $P_{\rm orb}$ in days and π in [mas].

3 ATMOSPHERIC PARAMETERS AND SPECTRUM SYNTHESIS

Relevant information about our program stars is summarized in Table 2. The $uvby\beta$ indices (de-reddened using the UVBYBETA code of Moon & Dworetsky 1985) were taken from Renson (1991). Geneva and UBV photometry were from Mermilliod, Mermilliod & Hauck (1997). The improved Hipparcos parallaxes were taken from van Leeuwen (2007). Table 2 also lists the absolute M_V magnitudes obtained from these parallaxes and V photometry. The atmospheric parameters were derived from both $uvby\beta$ and Geneva photometry. If both estimates were available we accepted their rounded mean as the best choice for model atmosphere parameters. All these stars seem to be SB1 binaries or have only a very weak secondary spectrum, hence the possible influence of their companions on photometry was neglected.

A detailed spectrum synthesis of the spectral regions was accomplished following the same recipe as in the previous papers of the series.

4 RESULTS FOR INDIVIDUAL STARS

The abundances obtained by synthetic spectrum fitting analysis are expressed relative to the Sun in terms of $[\mathrm{N/H}] = \log(\mathrm{N/H})_\star - \log(\mathrm{N/H})_\odot$ in the Table 3. Taking into account the accuracy of the atmospheric parameters, as well as the atomic data, the abundances of Al, Si, S, Ca and Fe are generally determined within $\lesssim 0.2$ dex, while the abundances of the other elements, which mainly occur in weak blends, are only approximate. The Ba abundances should be used with caution as they are usually derived from only one line (Ba II 6497 Å), which is at the edge of our frames. Apart from abundances and atmospheric parameters, we also derived the basic stellar properties like mass and age. We used the determined absolute magnitudes, created an HR diagram, and interpolated the evolutionary tracks and isochrones of Lejeune & Schaerer (2001). The masses, ages, and expected terminal-age main sequence (TAMS) obtained are also listed in Table 3 and the position of our programme stars in the HR diagram is illustrated in Figure 1. Both the synthetic and observed spectra are depicted in Figure 3.

Radial velocities, projected rotational velocities and microturbulent velocities determined as by-products are listed also in the Tables 1 and 3. A comparison between mea-



Figure 2. Measured radial velocities as listed in Table 1 (dots) in comparison with the predicted radial velocity curves (solid lines) versus phase. Orbital elements are taken from the appropriate references shown in Table 2.



Figure 3. Synthetic (solid) and observed (dashed) spectra of all of our programme stars. The observed spectra were shifted to the laboratory frame of synthetic spectra. A proper shift to the vertical coordinate was applied to all but the lowest spectrum.

sured radial velocities of the six stars and predicted radial velocity curves is shown in Figure 2. For the Am stars with $v \sin i < 50$ km s^{-1} the radial velocities of the primary stars were measured using the cross-correlation of the whole spectral region of the observed spectrum with the synthetic spectra. For the fast rotating star (HD 116657) with $v \sin i > 50$ km s^{-1} the cross-correlation technique produces large errors due to heavy line blending in some spectral regions. Consequently, in the Li-region, we restricted the cross-correlation region to 6710–6765 ÅÅ, and in the Ca region, we measured the velocities from the Ca I 6439 Å line using the center of mass method. The measurements of Budaj & Iliev (2003) with the same telescope configuration and center of mass method demonstrated, on the example of the fast rotating star (HD 178449, $v \sin i = 139$ km s^{-1}), that the standard deviation of such radial velocity measurements was less than 2 km s^{-1}. A discussion of the individual stars follows.

4.1 HD 116657

HD 116657 (ζ UMa B, HR 5055, ADS 8891 B, BD +55 1598B, SAO 28738, CSV 101382, A1m) is a part of the Mizar system. This system has been the first double star observed

Table 3. Abundances derived in terms of [N/H] for our six stars. Abundances of the Sun are in terms of $\log(N_{\rm el}/N_{\rm H}) + 12.00$

	Sun	HD 116657	HD 138213	HD 155375	HD 159560	HD 196544	HD 204188
Li	1.10	$\leq +2.08$	$\leq +2.2$	$\leq +1.88$	+1.68	$\leq +2.4$	$\leq +0.6$
C	8.52	≤ -0.62	≤ -0.18	≤ -0.11	-0.57	≤ -0.34	-0.82
O	8.83	≤ -0.38	-0.13	≤ -0.22	≤ -0.16	≤ -0.29	-0.13
Al	6.47	+0.30	–	–	–	–	–
Si	7.55	+0.00	+0.34	+0.05	+0.01	+0.10	+0.04
S	7.33	+0.10	+0.19	+0.0	+0.03	+0.15	-0.31
Ca	6.36	-0.42	+0.11	-0.64	-0.76	-0.50	-0.18
Ti	5.02	+0.00	+0.28	+0.14	-0.04	+0.24	+0.21
Fe	7.50	+0.34	+0.29	+0.22	+0.29	+0.27	+0.07
Ni	6.25	+0.67	+0.40	+0.65	+0.51	–	+0.23
Ba	2.21	–	+2.15	+1.85	+1.64	+1.19	+1.41
$\xi_{\rm turb}$	–	2.0	2.0	2.1	2.7	2.4	2.0
$v\sin i$	–	51	32	31	42	43	36
M	–	2.10	2.49	2.26	1.62	2.20	1.67
$\log T$	–	8.72	8.43	8.05	8.96	7.43	6.5*
$\log TAMS$	–	8.77	8.8	8.99	9.32	9.02	9.28

Note: Sun – abundances are taken from Grevesse & Sauval (1998) (recall that normal lithium abundance in hot stars or meteorites is [Li/H]=2.00); microturbulence - $\xi_{\rm turb}$ and $v\sin i$ are in km s^{-1}, M is mass in $M_{\odot}$, the age T and the Terminal-Age-Main-Sequence TAMS are given in years; * - the age is only upper limit (see details in the text)



Figure 4. Observed spectra of HD 155375. A proper shift to the vertical coordinate was applied to all but the lowest spectrum. Spectra with low S/N are not shown in the figure.

photographically. HD 116657 is the fainter star of the system where both stars are spectroscopic binaries.

The first spectral classification was given by Roman (1949) - A2/A8/A7 (Ca II/H/metal lines). Later Cowley et al. (1969) determined the spectral class as A1 from K-lines and Levato & Abt (1978) as A2/A7/A9 from K/H/metallic lines. Finally Abt & Levy (1985) determined the spectral class of HD 116657 as A1/A4/A3 according to K/H/metallic lines. The orbital elements of the binary system were obtained by Gutmann (1965) - $P_{\rm orb} = 175.6^d$, $K = 6.4$ km s^{-1}, e=0.463, $V_0 = -9.3$ km s^{-1}, $\omega = 6^{\circ}.9$. There have been numerous evaluations of the projected rotational velocity given by different authors. Meadows (1961) and Abt & Moyd (1973) measured $v\sin i = 50$ km s^{-1}. Dobrichev (1985) gave close value - $v\sin i = 52$ km s^{-1} while Slettebak (1954) obtained higher velocity - $v\sin i = 75$ km s^{-1}. Abt & Morrell (1995) measured the projected rotational velocity as $v\sin i = 51$ km s^{-1} but scaling this value to the results of Royer et al. (2002) changed the velocity to $v\sin i = 61$ km s^{-1}. Monier

(2005) obtained also $v\sin i = 61$ km s^{-1}. Our value of $v\sin i = 51$ km s^{-1} is very close to the values given by the majority of the authors. A few evaluations of the effective temperatures have been noted in the literature. The first value given by Cayrel de Strobel (1960) determined HD 116657 as a very cool star - $T_{\rm eff} = 5130$ K. The later values have given higher temperatures for the star. According to Smith (1971) the effective temperature was $T_{\rm eff} = 8800$ K and according to King et al. (2003) and Monier (2005) - $T_{\rm eff}$=8425 K. Our value of $T_{\rm eff}$=8470 K is very close to these authors' value.

The abundances of Ca, Fe, S are determined with good precision. For lithium we can only set an upper limit and claim that Li is not overabundant relative to the cosmic Li abundance. However, there is an indication of Li line. If this is confirmed then Iliev et al. (1998) Li abundance would be underestimated. The values listed for C and O are only upper limits also. According to our analysis Al is overabundant and Si seems normal from only one line SiI λ6721 Å . Ti could be evaluated only approximately to be normal from two blends of CaI λ6717 Å and FeI λ6678 Å. Ni is found overabundant. Some FeII lines are stronger than expected and it might indicate not only higher microturbulence or effective temperature but also lower gravity.

4.2 HD 138213

HD 138213(HR 5752, HIP 75770, BD + 47 2227) is a spectroscopic binary star. It was classified for the first time as a marginal metallic line star A5m: by Cowley et al. (1969). Later Eggen (1976) determined HD 138213 as a possible Am star based on photometry and Floquet (1975) suggested that the star was spectroscopically variable. According to Abt & Morrell (1995) the star was A2 IV. The orbital elements were taken by Lucy & Sweeney (1971) - $P_{\rm orb} = 105.95^d$, $K = 10.8$ km s^{-1}, e=0, $V_0 = -17.10$ km s^{-1}, $\omega = 0°$. Our radial velocities do not agree very well with the predicted velocity curve. It might be due to a small phase shift which accumulated over the years. There have been a few evaluations of the projected rotational velocity. Abt (1975) gave $v\sin i = 30$ km s^{-1} but later corrected it to $v\sin i = 45$ km s^{-1} (Abt & Morrell (1995)). Royer et al. (2002) scaled this value to their system and determined the velocity as $v\sin i = 54$ km s^{-1}. Our value is closer to the result given by Abt (1975) - $v\sin i = 32$ km s^{-1}.

Our analysis confirms that HD 138213 is a marginal Am star - Fe is overabundant and O is underabundant. Ca is almost solar abundant. For C and Li we give only the upper limits.

4.3 HD 155375

HD 155375(HR 6385, BD +12 3161, HIP 84036, SAO 102632, A1m) is a spectroscopic binary star. Osawa (1958) determined the spectral class of the star as A1/A3/A5 from K/H/metallic lines and Cowley et al. (1969) - as A1. Later Abt & Morrell (1995) specified it as A2 III class and Paunzen et al. (2001) - as A3 V class. Abt & Morrell (1995) also determined the projected rotational velocity as $v\sin i = 25$ km s^{-1} but scaling this value to the system of Royer et al. (2002) the latter changed the velocity to $v\sin i = 33$ km s^{-1}. Our value of $v\sin i = 31$ km s^{-1} is in good agreement with Royer et al. (2002) result. We used the orbital elements given by Debernardi (2002): $P_{\rm orb} = 23.25^d$, $K = 31.42$ km s^{-1}, e=0.422, $V_0 = 1.03$ km s^{-1}, $\omega = 114°.86$. The radial velocities measured from our spectra are in excelent agreement with the radial velocities calculated by using these elements (see Figure 2).

There are many Fe lines in the spectral region of 6400-6500 ÅÅ , so the abundance of Fe is very well determined. Ca is underabundant. The abundances given at Table 3 for C and O are upper limits. The situation with Li is the same as in the case of HD 116657 - the Li line is very weak and the obtained Li abundance is only upper limit.

More than one spectrum have been obtained in order to check the possible variability of some lines in the spectrum of HD 155375. As it is seen, a few lines have changed their profiles (see Figure 4). The relative changes of two lines, FeI λ6419.95 Å and FeI λ6421.35 Å, have been most obvious. Two calcium lines, CaI λ6439.08 Å and CaI λ6462.57 Å, have shown changes, too. The center of the lines has been changed and also there could be seen some features emerging from the blue side of the lines. All these observable clues forced us to suspect HD 155375 as a new SB2 star.

4.4 HD 159560

HD 159560 (ν^2 Dra, HR 6555, BD +55 1945, HIP 85829, SAO 30450, ADS 10628 A, A4m) is a member of the visual binary system. The angular separation between the components is 61.9″. Both components of the binary system are Am stars. There have been many determinations of the stellar spectral class in the literature. The first evaluation was given by Slettebak (1949) - A2/F0/F5 IV from K/H/metallic lines. Later the author specified the spectral class from H-lines as A7 (Slettebak 1963). According to Abt & Cardona (1984) the star was A4/F2V/F3. Cowley et al. (1969) also determined the spectral class of HD 159560 as A4 from H-lines. The evaluations of the projected rotational velocity of the star have been very different. Abt & Moyd (1973) gave $v\sin i = 35$ km s^{-1}, Böhm-Vitense & Dettmann (1980) - $v\sin i = 47$ km s^{-1}, Abt & Levy (1985) - $v\sin i = 50$ km s^{-1}, Abt & Morrell (1995) - $v\sin i = 58$ km s^{-1}. Finally, Royer et al. (2002) determined the rotational velocity of the star as $v\sin i = 68$ km s^{-1} scaled from the result of Abt & Morrell (1995). We obtained $v\sin i = 42$ km s^{-1} and this is the value of the rotational velocity we used for spectrum synthesis. We used the two different set of orbital elements given by Abt & Levy (1985) and Margoni et al. (1992) in order to check our radial velocities. Our results are in good agreement with the radial velocity curve obtained by using the orbital elements of the latter authors (see Figure 2). They gave the following orbital elements - $P_{\rm orb} = 38.034^d$, $K = 10.0$ km s^{-1}, e=0.03, $V_0 = -16.0$ km s^{-1}, $\omega = 92°$.

Our results of the abundances define the star as Am star. The line of Li I λ 6707 Å is well identified and is relatively stronger comparing to the other stars in this investigation. That is why the obtained abundance of Li is well determined.

4.5 HD 196544

HD 196544 (ι Del, HR 7883, BD +10 4339, HIP 101800, SAO 106322) is an A2V spectroscopic binary star. Adams (1912) determined for the first time the stellar radial velocity and obtained that it was variable. He classified the star as A2 spectral type star. Later Osawa (1959) determined the spectral class as A2 V according to MK system and as A4 from metallic lines. Levato (1975) confirmed the spectral class as A2 V and obtained the projected rotational velocity $v\sin i = 55$ km s^{-1} but later revised the velocity to $v\sin i = 60$ km s^{-1} (Garcia & Levato 1984). On the other hand, Abt & Morrell (1995) classified the star as A1 IV and gave a smaller value of the rotational velocity - $v\sin i = 30$ km s^{-1}. Royer et al. (2002) derived the projected rotational velocity and after merging it with the data available obtained $v\sin i = 41$ km s^{-1}. The orbital elements were derived by Harper (1935) - $P_{\rm orb} = 11.039^d$, $K = 26.0$ km s^{-1}, e=0.23, $V_0 = -4.9$ km s^{-1}, $\omega = 61^\circ.8$. Again, as in the case of HD 138213 the differences between our radial velocity measurements and the predicted velocity curve are due to a small phase shift accumulated over the years.

There have been a few elements abundances published in the literature. Lemke (1989), Lemke (1990) and Rentzsch-Holm (1997) gave the abundances of Fe, Ti, C, Ba, N and S using the atmospheric parameters $T_{\rm eff}$=9100 K and log g=4.3 which were very close to our values (see Table 2).

According to the abundances obtained by us, HD 196544 seems to be an Am star. The abundances of Fe and Ca are very well determined as Ca is underabundant and Fe is overabundant. Li line of 6707 Å is very weak and the abundance of Li is determined as only an upper limit. Other elements like C, O and Ti have weak lines in this spectral region, so their abundances should be scrutinized as upper limits. The differences between the abundances of Fe, Ti and C obtained by us and published by Lemke (1989) are within the errors.



Figure 5. [Ca/Fe] versus the eccentricity of the orbit. A least-squares regression line and the 95 percent confidence limits are drawn. The correlation coefficient is -0.55±0.05. HD 198391 is denoted by an asterisk. Two non-Am stars HD 178449 and HD 18778 denoted by open circles are given only for completeness and not included in the analysis.



Figure 6. [Ca/Fe] versus the orbital period. The periods are presented in logarithmic scale for more clarity. The correlation coefficient is -0.33±0.06.

The symbols are the same as in the previous figure.

4.6 HD 204188

HD 204188(IK Peg, HR 8210, HIP 105860, BD +18 4794, WD 2124+191, A8m) is an interesting single-lined spectroscopic binary with a companion star which is a massive white dwarf. Cowley et al. (1969) identified HR 8210 as a marginal Am star and determined the spectral class as A8m: but Abt & Bidelman (1969) identified it as a definite Am star. According to Bertaud (1970) the spectral class of HR 8210 was between A5 and F0. Later Guthrie (1987) in his study of the calcium abundances in metallic-line stars found that Ca was almost solar abundant. The most completed spectral identification was made by Abt & Morrell (1995) - A6/A9/F0 from K/H/metallic lines. Kurtz (1978) obtained that the primary Am star was also δ Sct star. Till now there are only a few stars which combine in one and the same star such contradictory characteristics. The first orbit determination was made by Harper (1927) who found the period of about 27 days and almost circular orbit. Later he clarified the period (Harper (1935)) and Batten, Fletcher & Mann (1978) assumed the eccentricity as e=0. We used the orbital parameters from the Ninth Catalogue of Spectroscopic Binary Orbits (SB9)(Pourbaix et al. (2004): $P_{\rm orb} = 21.724^d$, $K = 41.5$ km s^{-1}, e=0, $V_0 = -12.4$ km s^{-1}, $\omega = 0^\circ$. There have been many evaluations of the projected rotational velocity of IK Peg in the literature - from $v\sin i = 80$ km s^{-1} (Levato (1975)) to $v\sin i = 31$ km s^{-1} (Abt & Morrell (1995)). Royer et al. (2002) determined the projected rotational velocity as $v\sin i = 40$ km s^{-1} scaling the results of Abt & Morrell (1995). The projected rotational velocity obtained by us $v\sin i = 36$ km s^{-1} is in the range of Abt & Morrell (1995) and Royer et al. (2002) and very close to the values given by Rodriguez, Lopez-Gonzalez & Lopez de Coca (2000).

The iron-peak elements like Fe, Ti and Ni are sligthly overabundant and those of Ca and O - underabundant. For Li we present only an upper limit. As for the other stars of this study Ba is overabundant. These results as well as the atmospheric parameters are in good agreement with the results of Smalley et al. (1996). Our abundance analysis confirms that HD 204188 is a mild metallic line star.

The age obtained for this star is determined with not very high accuracy. We agree with Künzli & North (1998) who noticed the difficulty of determing the stellar age properly when the star is near ZAMS.

5 TIDAL, ROTATION AND EVOLUTION EFFECTS IN AM PHENOMENA

Following the systematic search for abundance anomalies of Am stars driven by tidal effects we continued with studying the dependences between the chemical abundances and the orbital elements of the binary systems, the projected rotational velocity and the physical parameters of Am stars like mass, age, temperature. The main advantage of our analysis is the homogeneity of the observational material we used - all data were obtained at one telescope with the same spectrograph and detector as well as processed with the same data reduction package and analysed with the same code Synspec.

As it is well known Am peculiarities are mainly manifested by Ca deficit and Fe overabundances and they could be represented in a reliable way through the values of [Ca/Fe] where [Ca/Fe]=[Ca/H]-[Fe/H]. This ratio would multiply the effect of Am peculiarities because these two elements have shown an opposite behaviour. Besides the lines of both elements get weaker with the increasing temperature and consequently [Ca/Fe] ratio is not that sensitive to the uncertainties of the effective temperature.

Up to now fifteen stars from our sample were fully processed. The results are taken from this work and also from Paper I and Paper II. Two of the stars in this sample, HD 178449 and HD 18778 were found not to be Am stars. They are put on the figures but not included in the analysis.

First we investigate the dependences of [Ca/Fe] from the eccentricity (see Figure 5). Despite of some scatter in the data there seems to be a trend like that mentioned earlier by Budaj (1997) and Iliev et al. (1998). The correlation coefficient is -0.55±0.05. One star, HD 198391 denoted by an asterisk in the figures, is distinguished from the common trend. This star is the hottest star amongst the sample and it is not a typical Am star (Paper I). The star could be a transitional object between Am and the hotter HgMn stars because of its position in HRD close to the region occupied by HgMn stars and also of its abundances (for more details see Paper I). Namely, that Am peculiarities increase ([Ca/Fe] decreases) with increasing eccentricity of the binary system. The dependence of [Ca/Fe] from the orbital period shown in Figure 6 is not so clear but still there is a tendency the metalicity to increase towards the longer periods. Of course, these two parameters characterizing the binary system, eccentricity and period, are not fully independent because of the synchronization and circularization of the orbits. This was the main reason to analyse only stars with periods between 10 and 200 days. The comparison of the three parameters simultaneously in 3D-graph (see Figure 7) - metalicity, eccentricity and period showed us that there was not any correlation between period and eccentricity in the whole range of periods which extends even beyond 150 days.



Figure 7. The dependence between [Ca/Fe], the eccentricity and the orbital period. Open circles stand for plane projections.

We study also the dependence of the Am peculiarities on the projected rotational velocity of the Am stars (see Figure 9). In general, there is a clear trend of increasing the peculiarity (decreasing of [Ca/Fe]) towards small values of $v \sin i$ - the correlation coefficient is +0.85±0.04. Besides the hottest star already mentioned, HD 198391, two other stars, HD 138213 and HD 204188, both marginal Am stars, do not follow the common trend. The fact that these two stars depart from the clear smooth correlation of metallicity and $v \sin i$ indicates that the rotation is not the only agent responsible for this peculiarity. It confirms our claims from Iliev et al. (1998) that the low Am peculiarity in these two systems is due to their small eccentricity. Again in order to check the possible correlation between the parameters studied we tried to combine [Ca/Fe], the eccentricity and $v \sin i$ in 3D-graph (see Figure 8). We obtained a weak negative correlation between the eccentricity and the projected rotational velocity - faster rotating stars tended to have smaller eccentricities.

The dependence of the Am peculiarity on the effective temperature is plotted on the Fig. 10. There might be a trend of decreasing the peculiarity with the temperature - the correlation coefficient is +0.36±0.06.

We sought a possible dependence of the Am peculiarity on the age and the mass of the stars. With the exception of one star, HD 196544, the other stars studied have very close ages - the difference between the youngest and the oldest star is 0.51 in $\log T$ (where T is the age in years). The majority of the masses of the stars is also in very short interval - 0.5 in solar masses. As a result from all our data we can not see any signs of dependence between these parameters (mass and age) and the Am peculiarity.

We also studied the connection between the micro-



Figure 8. The dependence between [Ca/Fe], the eccentricity and the projected rotational velocity. Again, open circles stand for plane projections.



Figure 9. [Ca/Fe] versus the projected rotational velocity. The correlation coefficient is +0.85±0.04. The symbols are the same as in Figure 5.

turbulence and the effective temperature (see Fig. 11). There seems not to be any dependence between these parameters for the temperature region 7000-9000 K. But at higher temperatures it is possible to claim that the microturbulence decreases with the increasing temperature. These results fit relatively well with the conclusion given by Burkhart & Coupry (1992). The microturbulence is a pure fitting parameter which brings into agreement the abundances from weak and strong lines. It does not necessarily have the meaning of existing turbulent motions. Nevertheless, this behaviour is in agreement with the expectations and with the fact that supeficial convective zones get thinner at higher temperatures and cease at about 10 000 K. We also studied a possible dependence of the Am peculiarities on the microturbulence but there is no clear correlation. Apparently, the microturbulence does not seem to destroy the Am peculiarity.



Figure 10. [Ca/Fe] versus the effective temperature. The correlation coefficient is +0.36±0.06. The symbols are the same as in Figure 5.



Figure 11. The dependence of the microturbulence from the effective temperature. The symbols are the same as in Figure 5.

We compared our measurements of the projected rotational velocities of the program stars to those obtained by Abt & Morrell (1995) and by Royer et al. (2002). As it is seen from Figure 12 our measurements are in good agreement with them. The velocities of Abt & Morrell (1995) seems to be closer to our values while those of Royer et al. (2002) are often slightly higher.



Figure 12. The projected rotational velocities obtained by us versus those given by Abt & Morrell (1995) (open circles) and by Royer et al. (2002) (filled circles).

6 CONCLUSIONS

In this third and last paper, we have analysed another six binaries from our sample and derived their chemical composition, temperatures, gravities, projected rotational velocities, masses and ages. The obtained abundances of certain elements allowed us to conclude that these stars are Am stars. We suggested that one of them, HD 155375 could be a new SB2 stars based on 5 spectra obtained. The tidal interaction in all binary systems studied by us till now was explored. There is clear dependence of the Am peculiarity on the projected rotational velocity for clear defined Am stars. Apart from that there seems to be a correlation of the Am peculiarity with the eccentricity and may be also with the orbital period and a weak anticorrelation with the effective temperature.

Assuming that the Am peculiarity is due to the microscopic diffusion processes in stable atmospheres then this dependence of the Am peculiarity on the eccentricity must be due to some mechanism which will stabilize the atmosphere of the star and reduce mixing.

One could speculate that star on an eccentric orbit is subject to a variable gravitational perturbance or oscillations. These might cause departures from the single star rotation, drive the star towards the pseudo-synchronisation, and might reduce the differential rotation and/or rotationally induced mixing.

This confirms the hypothesis of Budaj (1996,1997) about the existence of a stabilization mechanism due to the companion of the star and that this mechanism operates up to the orbital period of about 200 days and that it is relevant for the Am phenomenon.

ACKNOWLEDGMENTS

IS and II acknowledge the partial support from Bulgarian NSF under grant D0 02-85. JB gratefully acknowledges grant support from the Marie Curie International Reintegration Grant FP7-200297.

This research was supported partially by the VEGA grants 2/0074/09, 2/0078/10 and 2/0094/11 from the Slovak Academy of Sciences. This study made use of IRAF Data Reduction and Analysis System and the Vienna Atomic Line Data Base (VALD) services.

We wish also to thank the referee P. North for his valuable comments and suggestions that helped us to improve substantially the paper.

REFERENCES

Abt H.A., 1975, ApJ, 195, 405
Abt H.A., 2000, ApJ, 544, 933
Abt H.A., Bidelman W., 1969, ApJ., 158, 1091
Abt H.A., Cardona O., 1984, ApJ, 276, 266
Abt H.A., Levy S.G., 1985, ApJS, 59, 229
Abt H.A., Moyd K.I., 1973 ApJ, 182, 809
Abt H.A., Morrell N.I., 1995, ApJS, 99, 135
Adams W.S., 1912, ApJ, 35, 163
Batten A.H., Fletcher J.M., Mann P.J., 1978, PDAO 15, 121
Bertaud C., 1970, A&AS, 1, 7
Boffin, H. M. J., 2010, A & A, 524, 14
Böhm-Vitense E., Dettmann T., 1980, ApJ, 236, 560
Budaj J., 1996, A&A, 313, 523
Budaj J., 1997, A&A, 326, 655
Budaj J., Iliev I.Kh., 2003, MNRAS, 346, 27 (Paper I)
Burkhart C., Coupry M.F., 1992, A&As, 95, 41
Burkhart C., Coupry M.F., 2000, A&A, 354, 216
Carquillat J.-M., Prieur J.-L., 2007, MNRAS, 380, 1064
Cayrel de Strobel G., 1960, AnAp, 23, 278
Cowley A., Cowley C., Jaschek M., Jaschek C., 1969, AJ, 74, 375
Debernardi Y., 2002, PhD Thesis
Debernardi Y., Mermilliod J.-C., Carquillat J.-M., Ginestet N., 2000, A&A, 354, 881
Dobrichev V., 1985, AISof, 4, 40
Eggen O.J., 1976, PASP 88, 402
ESA, 1997, in The Hipparcos and Tycho Catalogues, ESA SP-1200
Fekel Francis C., Williamson Michael H., 2010, AJ, 140, 1381
Floquet M., 1975, A&AS, 21, 25
Fossati L., Bagnulo S., Monier R., Khan S.A., Kochukhov O., Landstreet J., Wade G., Weiss W., 2007, A & A, 476, 911
Garcia B.E., Levato H., 1984, RMxAA, 9, 9
Grevesse N., Sauval A.J., 1998, SSRv, 85, 161
Guthrie B.N.G., 1987, MNRAS, 226, 361
Gutmann F., 1965, PDAO, 12, 361
Harper W.H., 1927, PDAO, 4, 161
Harper W.H., 1935, PDAO, 6, 207
Hubrig S., González J., Schöller M., 2010, ASPC, 435, 257
Hui-Bon-Hoa A., 2000, A&AS, 144, 203
Iliev I.Kh., Budaj J., 2008, Contrib. Astron. Obs. Skalnate Pleso, 38, 134
Iliev I.Kh., Budaj J., Feňovčík M., Stateva I., Richards M.T., 2006, MNRAS, 370, 819 (Paper II)
Iliev I.Kh., Budaj J., Zverko J., Barzova I.S., Žižňovský J., 1998, A&AS, 128, 497
King J.R., Villareal A.R., Soderblom D.R¿, Gulliver A.F., Adelman S.J., 2003, AJ, 125, 1980
Künzli M., North P., 1998, A&A, 330, 651

Kurtz D.W., 1978, ApJ, 221, 869

Lejeune T., Schaerer D., 2001, A&A, 366, 538

Lemke M., 1989, A&A, 225, 125

Lemke M., 1990, A&A, 240, 331

Levato H., 1975, A&AS, 19, 91

Levato H., 1975, A&AS, 19, 91

Lucy L.B., Sweeney M.A., 1971, AJ, 76, 544

Margoni R., Munari U., Stagni R., 1992, A&AS, 93, 545

Meadows A.J., 1961, ApJ, 133, 907

Mermilliod J.-C., Mermilliod M., Hauck B., 1997, A&AS, 124, 349

Mikulášek Z., Zverko, J.; Žižňovský J., Krtička J., Iliev I.Kh., Kudryavtsev D. O., Grf T., Zejda M., 2010, ASPC, 435, 319

Monier R., 2005, A&A, 442, 563

Monier R., Richard O., 2004, in Zverko J., Žižňovský J., Adelman S.J., Weiss W.W., eds, IAU Symp., No. 224., The A-Star Puzzle. Cambridge University Press, Cambridge, UK, p. 209

Moon T. T., Dworetsky M. M., 1985, MNRAS, 217, 305

North P., Ginestet N., Carquillat J.-M., Carrier F., Udry S., 1998, CAOSP, 27, 179

Osawa K., 1958, PASJ, 10, 102

Osawa K., 1958, ApJ, 130, 159

Paunzen E., Duffee B., Heiter U., Kuschnig R., Weiss W.W., 2001, A&A, 373, 625

Pourbaix D., Tokovinin A.A., Batten A.H., Fekel F.C., Hartkopf W.I., Levato H., Morrell N.I., Torres G., Udry S., 2004, A&A, 424, 727

Quiroga C., Torres A. F., Cidale L. S., 2010, A & A, 521, 75

Renson P., 1991, Catalogue Général des Etoiles Ap et Am (Astr. Inst.-Univ. de Liège)

Rentzsch-Holm I., 1997, A&A, 317, 178

Rodriguez E., Lopez-Gonzalez M.J., Lopez de Coca P., 2000, A&AS, 144, 469

Roman N.G., 1949, ApJ, 110, 205

Royer F., Grenier S., Baylac M.-O., Gómez A.E., Zorec J., 2002, A&A, 393, 897

Slettebak A., 1949, ApJ, 109, 547

Slettebak A., 1954, ApJ, 119, 146

Slettebak A., 1963, ApJ, 138, 118

Smalley B., Smith K.C., Wonnacott D., Allen C.S., 1996, MNRAS, 278, 688

Smith M., 1971, A&A, 11, 325

van Leeuwen F., 2007, A&A, 474, 653

Zhao M., Monnier J. D., Torres G., Boden A. F., Claret A., Millan-Gabet R., Pedretti E., Berger J.-P., Traub W. A., Schloerb F. P., Carleton N. P., Kern P., Lacasse M. G., Malbet F., Perraut K., 2007, ApJ, 659, 626

Zverko J., Žižňovský J., Mikulášek Z., Iliev I. Kh., 2008, Contrib. Astron. Obs. Skalnate Pleso, 38, 467

Zverko J., Žižňovský J., Mikulášek Z., Krtička J., Iliev I. Kh., Stateva I.K., Romanyuk I. I., Kudryavtsev D. O., 2009, A&A, 506, 845